



# Achillea

## SASSY SUMMER



Cuttings

Achillea, also known as Yarrow, was named after the ancient Greek hero Achilles. According to the legend, Achilles used Yarrow to treat his and his soldier's wounds during the Trojan War.

Achillea is a symbol of healing and protection.

Sassy Summer is a taller, more substantial presence than any other series in the market.

They have a taller height, all members have sturdy flower stems that will look great at each bouquet or arrangement.

With a range of beautiful colors, such as yellow, red, pink, and orange, you'll find something that fits your needs.

SUITE:



Taffy



Lemon



Sangria



Sunset

TECHNICAL INFORMATION

CROP TIME

From planting to harvest: 14 weeks

After the first flush it produces again 8 to 10 weeks later for 8-9 weeks.

PLANTING DENSITY

6 plants/ m<sup>2</sup> net

PRE-PLANTING

**Soil:** Well drained

**Netting:** 2 levels

pH: 5.5 – 6.2

TEMPERATURE

Ideal for growth: 15-18 °C

PINCH

If the plant is not well developed with a 25 to 30 cm diameter, pinch the stems produced by the plant (pinch at ground level).

Once reached the desired size, apply 50 ppm of gibberellic acid. Then stop the pinching and let them grow. Once the production cycle finishes, wait until the plant has had enough size to reapply gibberellic acid.

IRRIGATION

Avoid excess of humidity. The achillea is tolerant to dry conditions.

FERTILIZATION

Doesn't require high fertilization. A general fertilization formula can be the following::

N: 80 ppm , P: 50 ppm, K: 100 ppm

Ca: 100 ppm, Mg: 40 ppm, S: 50 ppm

Fe: 2 ppm, Mn: 2.5 ppm, Cu: 0.2 ppm

Zn: 0.2 ppm, B: 0.2 ppm.

HARVEST

Harvest with flowers fully opened where pollen is seen. Cut the stems at ground level without leaving stumps.

POST-HARVEST

Hydrate in Chrysal L&A: 10 gr /lt for 6 hours.

PESTS & DISEASES

- Aphids
- Rhizoctonia
- Leaf miner
- Thrips